

Information about the scientific activities of the Cesa

Free irregular online issues of the Cesa News are regularly archived in the recognized "Internet Archive", in accordance with the Rules of the ICZN

Announcements- General News – Expeditions-
Publications –Visitors – Workshops - Seminars -
&c

Nr.149

28 November 2017

Research Article

<http://zoobank.org/urn:lsid:zoobank.org:pub:C07D3129-F21F-4BC7-98B9-1C76BE9A4266>

Preliminary list of the Lepidoptera recorded from Erek Mountain (Van Province, East Turkey)

Muhabbet Kemal ¹ Ahmet Ömer Koçak

Abstract: Preliminary list of the moths recorded from Erek Mountain (Van Province, East Turkey). *Cesa News* 149: 1-15, 37 figs.

In this paper, 196 species of 17 families are listed from Erek Mt (Van Province, East Turkey). Observational images of some species and genitalia used for identifications are added. Phenological and faunistical notes are given to some species. Early stage and larval food-plant of *Polygona egea* are mentioned and illustrated.

Keywords: Lepidoptera, fauna, Erek Mountain, Van, Turkey

Within the Project of the Cesa, announced recently in the ResearchGate "[Lepidoptera fauna of Erek Mountain \(Van Province, East Turkey\) \(LEM\)](#)", first list of the 196 lepidopteran species are given below. The species were collected by the authors from the mountain at irregular intervals. The butterflies of Erek Mountain in İpekyolu district (Van Province) was studied by Kayci (2007). His list has been updated today. The present paper deals with the butterfly observations and the moths recorded by the authors in the past from various places of Erek Mt., as well as some autumnal species collected during the past October (Fig. 17).

The authors listed 1179 Lepidopteran species known from Van, as well as the other pterygots of this province (Koçak & Kemal, 2012). Since then, the number of the species of Lepidoptera has reached to 1568, while the species of other pterygota 1477.

Erek (3200m) is a difficult, steep, and rocky mountain, especially from the standpoint of night collecting. Inside of the mountain, there is no road. On the other hand, the ring roads are very far from the mountain mass. By day, up to the elevation 2500m can be studied from the eastern slopes. But the night collecting in the next years will be probably restricted below 2300m. Under these circumstances, the expected number of the species of the Lepidoptera may not be so high. But yet, the fauna of this mountain is apparently under the influence of West Asiatic Elements zoogeographically. Therefore, interesting records are expected.

¹ Asst. Prof. Dr. Muhabbet Kemal, Van Yüzüncü Yıl University, Faculty of Science, Dept. of Biology, Campus, Van / Turkey.
e-mail: muhabbet_kemal@yahoo.com.tr

co-author: Prof. em. Dr. Ahmet Ömer Koçak, c/o Van Yüzüncü Yıl University, Faculty of Science, Dept. of Biology, Campus, Van / Turkey. e-mail: cesa_tr@yahoo.com.tr

Butterflies

Argynnidae

The family is represented by 13 species on the mountain.

1. *Aglais urticae* (Linnaeus, 1758)
2. *Argynnис (Fabriciana) niobe* (Linnaeus, 1758)
3. *Argynnис (Pandoriana) pandora* ([Denis & Schiffermüller], 1775)
4. *Brenthis hecate* ([Denis & Schiffermüller], 1775)
5. *Issoria lathonia* (Linnaeus, 1758)
6. *Melitaea (Cinclidia) phoebe* (Goeze, 1779)
7. *Melitaea (Didymaeformis) didyma* (Esper, [1779])
8. *Melitaea (Didymaeformis) persea* Kollar, [1849]
9. *Melitaea (Didymaeformis) syriaca* Rebel, 1905
10. *Melitaea (s.str.) cinxia* (Linnaeus, 1758)
11. *Polygonia (Comma) egea* (Cramer, [1775]) (Figs. 1-4)
12. *Vanessa (Cynthia) cardui* (Linnaeus, 1758)
13. *Vanessa (s.str.) atalanta* (Linnaeus, 1758)

Hesperiidae

The family is represented by 14 species on the mountain.

14. *Carcharodus (Lavatheria) stauderi* Reverdin, 1913
15. *Carcharodus (Reverdinus) orientalis* Reverdin, 1913
16. *Carcharodus (s.str.) alceaе* (Esper, [1780]) (Fig. 5)
17. *Eogenes alcides* Herrich-Schäffer, [1852]
18. *Hesperia comma* (Linnaeus, 1758)
19. *Muschampia poggei* (Lederer, 1858)
20. *Muschampia proteides* (F. Wagner, 1929)
21. *Pyrgus melotis* (Duponchel, [1834])
22. *Pyrgus serratulae* (Rambur, [1839])
23. *Pyrgus sidae* (Esper, [1784]) (Fig. 6)
24. *Spatialia (Neospatialia) orbifer* (Hübner, [1823])
25. *Spatialia (s.str.) phlomidis* (Herrich-Schäffer, [1845])
26. *Thymelicus lineolus* (Ochsenheimer, 1808)
27. *Thymelicus sylvestris* (Poda, 1761)

Lycaenidae

The family is represented by 40 species on the mountain.

28. *Callophrys danchenkoi* Zhdanko, 1998
29. *Callophrys mystaphia* Miller, [1913] (Kemal, 2009)
30. *Cupido (s.str.) osiris* (Meigen, [1829])
31. *Glaucoopsyche (s.str.) alexis* (Poda, 1761)
32. *Lycaena (Alciphronia) alciphron* (Rottemburg, 1775)
33. *Lycaena (Heodes) virgaureae* (Linnaeus, 1758)
34. *Lycaena (Palaeochrysophanus) candens* (Herrich-Schäffer, [1845])
35. *Lycaena (Thersamonia) asabinus* (Gerhard, [1850]) (Fig. 7)
36. *Lycaena (Thersamonia) kefersteinii* (Gerhard, [1850]) (Fig. 8)
37. *Lycaena (Thersamonia) kurdistanica* (Riley, 1921) (Figs. 9,10)
38. *Lycaena (Thersamonia) thetis* (Klug, 1834)
39. *Lycaena (s.str.) phlaeas* (Linnaeus, 1761) (Fig. 11)
40. *Plebejus (Kretania) carmon* (Gerhard, [1851])
41. *Plebejus (Plebejides) zephyrinus* (Christoph, 1884)
42. *Plebejus (s.str.) argus* (Linnaeus, 1758)
43. *Polyommatus (Albulina (Plebejidea)) loewii* (Zeller, 1847)
44. *Polyommatus (Albulina (Vacciniina)) alcedo* (Christoph, 1877)
45. *Polyommatus (Aricia (Ultraaricia)) crassipunctus* (Christoph, 1893)
46. *Polyommatus (Aricia (s.str.)) agestis* ([Denis & Schiffermüller], 1775)
47. *Polyommatus (s.str. (Agrodiaetus (Antidolus))) kurdistanicus* (Forster, 1961)
48. *Polyommatus (s.str. (Agrodiaetus (Antidolus))) menalcas* (Freyer, [1837])
49. *Polyommatus (s.str. (Agrodiaetus (Phyllisia))) vanensis* (De Lesse, 1957)
50. *Polyommatus (s.str. (Agrodiaetus (Transcaspius))) huberti* (Carbonell, 1993)

51. *Polyommatus (s.str. (Agrodiaetus (Transcaspius))) surakovi* Dantchenko & Lukhtanov, 1994
 52. *Polyommatus (s.str. (Agrodiaetus (Xerxesia))) cyaneus* (Staudinger, 1899)
 53. *Polyommatus (s.str. (Cyaniris)) bellis* (Freyer, [1842])
 54. *Polyommatus (s.str. (Lysandra)) bellargus* (Rottemburg, 1775)
 55. *Polyommatus (s.str. (Lysandra)) corydonius* (Herrich-Schäffer, [1852])
 56. *Polyommatus (s.str. (Meleageria)) daphnis* ([Denis & Schiffermüller], 1775)
 57. *Polyommatus (s.str. (Plebicula)) amandus* (Schneider, 1792)
 58. *Polyommatus (s.str. (Sublysandra)) aedon* (Christoph, 1877)
 59. *Polyommatus (s.str. (Thersitesia)) thersites* (Canterer, [1835])
 60. *Polyommatus (s.str.) icarus* (Rottemburg, 1775)
 61. *Pseudophilotes vicrama* (Moore, 1865)
 62. *Rubrapterus bavius* (Eversmann, 1832) (Fig. 12)
 63. *Satyrium (Armenia) ledereri* (Boisduval, 1848)
 64. *Satyrium (Nordmannia) abdominalis* (Gerhard, [1850])
 65. *Satyrium (Strymonidia) spini* (Fabricius, 1787)
 66. *Tomares callimachus* (Eversmann, 1848)
 67. *Turanana endymion* (Freyer, [1850])

Papilionidae

The family is represented by 4 species on the mountain.

68. *Archon apollinaris* (Staudinger, [1892])
 69. *Papilio (s.str.) machaon* Linnaeus, 1758
 70. *Parnassius (Driopa) mnemosyne* (Linnaeus, 1758)
 71. *Zerynthia (Allancastria) deyrollei* (Oberthür, 1869)

Pieridae

The family is represented by 16 species on the mountain.

72. *Anthocharis cardamines* (Linnaeus, 1758)
 73. *Anthocharis damone* Boisduval, 1836 (Fig. 13)
 74. *Anthocharis grueneri* Herrich-Schäffer, [1851] (Fig. 14)
 75. *Aporia (s.str.) crataegi* (Linnaeus, 1758)
 76. *Colias (Eriocolias) crocea* (Fourcroy, 1785)
 77. *Colias (Neocolias) aurorina* Herrich-Schäffer, [1850]
 78. *Gonepteryx (s.str.) rhamni* (Linnaeus, 1758)
 79. *Leptidea duponcheli* (Staudinger, 1871)
 80. *Pieris (Artogeia) ergane* (Geyer, [1828])
 81. *Pieris (Artogeia) krueperi* Staudinger, 1860
 82. *Pieris (Artogeia) rapae* (Linnaeus, 1758)
 83. *Pieris (s.str.) brassicae* (Linnaeus, 1758)
 84. *Pontia callidice* (Hübner, [1800])
 85. *Pontia chloridice* (Hübner, [1813])
 86. *Pontia edusa* (Fabricius, 1777)
 87. *Zegris eupheme* (Esper, [1804])

Satyridae

The family is represented by 22 species on the mountain.

88. *Arethusana arethusa* ([Denis & Schiffermüller], 1775)
 89. *Chazara (Neochazara) anthe* (Hoffmannsegg, 1804)
 90. *Chazara (s.str.) bischoffi* (Herrich-Schäffer, [1846])
 91. *Chazara (s.str.) briseis* (Linnaeus, 1764)
 92. *Coenonympha (s.str.) leander* (Fabricius, 1787)
 93. *Coenonympha (s.str.) pamphilus* (Linnaeus, 1758)
 94. *Esperarge (s.str.) clymene* (Fabricius, 1787)
 95. *Hyponephele (s.str. (Ereminephele)) naricoides* Gross, 1977
 96. *Hyponephele (s.str. (Tengrinephele)) cadusia* (Lederer, 1869)
 97. *Hyponephele (s.str.) lupina* (Costa, [1836])
 98. *Hyponephele (s.str.) lycaon* (Rottemburg, 1775)
 99. *Lasiommata maera* (Linnaeus, 1758) (Fig. 15)
 100. *Lasiommata megera* (Linnaeus, 1767)

101. *Maniola (Telmessiola) telmessia* (Zeller, 1847)
 102. *Maniola (s.str.) jurtina* (Linnaeus, 1758)
 103. *Melanargia (Turcargia) hylata* (Ménétriés, 1832)
 104. *Melanargia (Turcargia) syriaca* (Oberthür, 1894)
 105. *Pseudochazara (s.str.) beroe* (Freyer, [1843])
 106. *Pseudochazara (s.str.) geyeri* (Herrich-Schäffer, [1846])
 107. *Pseudochazara (s.str.) pelopea* (Klug, 1832)
 108. *Satyrus (Asatyrus (s.str.)) iranicus* Schwingenschuss, 1939
 109. *Satyrus (s.str.(s.str.)) favonius* Staudinger, [1892]

Moths

Mostly nocturnal. Only 87 species are reported here.

Arctiidae

Only 2 species have been recorded from Erek Mt. so far.

110. *Lacydes spectabilis* (Tauscher, 1806).

Locally common. Autumnal.

111. *Ocnogyna (s.str.) loewii* (Zeller, 1846)

Locally common. Autumnal.

Geometridae

Only 10 species have been recorded from Erek Mt. so far.

112. *Aplocera annexata* (Freyer, [1830])

(Fig.16)

Common in summer, diurnal & nocturnal.

113. *Catarhoe permixtaria* (Guenée, [1858])

Common in summer.

114. *Docirava mundulata* (Guenée, [1858])

Vernal, rare.

115. *Idaea rufaria* (Hübner, [1799])

Common in summer.

116. *Isturgia murinaria* ([Denis & Schiffermüller], 1775)

Vernal.

117. *Larentia clavaria* Haworth, [1809]

(Fig.18)

An autumnal species. Uncommon in Van Province. Previously recorded from Çatak (Saklıvadi) (Kemal & Koçak, 2017). They also attached a distributional map for this species.

118. *Lythria purpuraria* (Linnaeus, 1758)

Uncommon in summer.

119. *Rhodometra sacraria* (Linnaeus, 1767)

120. *Scopula immistaria* (Herrich-Schäffer, [1852])

121. *Scotopteryx kuznetzovi* (Vardikjan, 1957)

(Figs.19, 27-30)

Recently, Rajaei & Stüning (2012) reported this species from NW Iran and Turkey (Bitlis: Ahlat, Van: Güzeldere). The early stages, female and exact distribution of this autumnal species in Turkey are still unknown. Separately, it has been planned to research molecular data of this group. This species flies only in October-November in Van Province (Tuşba, İpekyolu and Gürpınar districts).

Hepialidae

122. *Triodia amasina* (Herrich-Schäffer, [1851])

Locally common. Crepuscular & nocturnal. It is usually attracted by the light in the early hours of the night. Autumnal.

Lasiocampidae

123. *Malacosoma castrensis* (Linnaeus, 1758)

(Fig.20)

Common in summer.

Lymantriidae

124. *Polymona lapidicola* (Herrich-Schäffer, [1852])

Common in summer.

Noctuidae

This species number of this family is the highest in Turkey. For the time being, 53 species have been recorded from Erek Mt. so far.

125. *Acontia (Emmelia) trabealis* (Scopoli, 1763)

Locally common.

126. *Agrochola (Alpichola) egorovi* (O. Bang-Haas, 1934) (Figs. 21, 31, 32)

For the identity of this species Ronkay (1984) is recommended. Autumnal.

127. *Agrotis (Crassagrotis) obesa* (Boisduval, 1829)

Autumnal.

128. *Agrotis (s.str.) desertorum* Boisduval, 1840

129. *Ammoconia anonyma* Ronkay & Varga, 1984

Autumnal.

130. *Amphipyra (s.str.) tragopoginis* (Clerck, 1759)

Uncommon.

131. *Behounekia freyeri* (Frivaldszky, 1835)

Vernal, common.

132. *Calymma communimacula* ([Denis & Schiffermüller], 1775)

133. *Caradrina (Paradrina) wulschlegeli* (Püngeler, 1903)

134. *Chersotis (s.str.) anatolica* (Draudt, 1936)

Autumnal.

135. *Chersotis (s.str.) laeta* (Rebel, 1904)

136. *Cosmia (Calymnia) trapezina* (Linnaeus, 1758)

137. *Cryphia (Bryoleuca) raptricula* ([Denis & Schiffermüller], 1775)

138. *Cryphia (Bryophila) maeonis* (Lederer, 1865)

139. *Cryphia (Bryophila) occidentalis* (Ostheder, 1933)

140. *Dichagyris (Albocosta) flammatra* ([Denis & Schiffermüller], 1775)

141. *Dichagyris (Yigoga) celsicola* (Bellier, 1858)

142. *Dichagyris (s.str.) forficula* (Eversmann, 1851)

143. *Dichagyris (s.str.) amoena* (Staudinger, 1892) (Fig.22)

Uncommon but widely distributed in East Turkey. Autumnal.

144. *Drasteria caucasica* (Kolenati, 1846)

145. *Drasteria saisani* (Staudinger, 1882)

146. *Earias clorana* (Linnaeus, 1761)

147. *Episema korsakovi* (Christoph, 1885)

Uncommon. Autumnal.

148. *Eublemma (Glossodice) polygrammum* (Duponchel, [1842])

149. *Eublemma (candidana-gr.) ochreolum* (Staudinger, 1900)

150. *Euchalcia emichi* (Rogenhofer, 1873)

151. *Euxoa (s.str.) homicida* (Staudinger, 1900)

152. *Euxoa (s.str.) scurrilis* Draudt, 1937

153. *Euxoa (s.str.) temera* (Hübner, [1808])

154. *Hadena (Anepia) syriaca* (Ostheder, 1933)

155. *Hadena (s.str.) compta* ([Denis & Schiffermüller], 1775)

156. *Hadena (s.str.) praetermissa* Hacker, 1996

157. *Hadena (s.str.) purpurea* Hacker, 1996

158. *Haemerosia renalis* (Hübner, [1813])

159. *Hecatera rhodocharis* (Brandt, 1938)

160. *Heptapotamia eustratii* Alpheraky, 1882 (Fig.23)

The early stage and larval food-plant were published by the authors (Kemal & Koçak, 2015). Local, and uncommon. Autumnal.

161. *Hoplodrina ambigua* ([Denis & Schiffermüller], 1775)

162. *Hoplodrina pfeifferi* (Boursin, 1932)

163. *Leucania loreyi* (Duponchel, 1827)

164. *Leucania punctosa* (Treitschke, 1825)

165. *Margelana flavidior* Wagner, 1831 (Fig.24)

Autumnal.

166. *Oligia latruncula* ([Denis & Schiffermüller], 1775)

167. *Opigena polygona* (Fabricius, 1787)

Locally common, in late summer and autumn.

168. *Phytometra viridaria* (Clerck, 1759)

169. *Polymixis (s.str.) csorbagabori* L.Ronkay, Varga & Hreblay, 1998 (Figs. 33, 34)

Autumnal, rare.

170. *Polymixis (s.str.) perchrysa* Hacker & L.Ronkay, 1992 (Figs. 25, 35)

This species was described from Ağrı Prov., Aydıntepe (Hacker & L.Ronkay, 1992: 488), and recently reported from Çatak, Saklıvadi (Van Prov.) (Kemal & Koçak, 2017). It is an autumnal, and apparently rare species in the region.

171. *Polymixis (s.str.) rosinae* (Bohatsch, 1910) (Figs. 36, 37)

Boursin (1944) described “*Antitype paradisiaca*” from Armenia, and illustrated the male genitalia. Boursin’s name is currently accepted as junior synonym of *rosinae* Bohatsch, 1910. We used the Boursin’s photograph of *paradisiaca* for identification of our specimen. All these *Polymixis* species mentioned here are found sympatric. Autumnal.

172. *Pseudenargia deleta* (Ostheder, 1933)

Autumnal.

173. *Rhypagla lacernaria* (Hübner, [1813])

174. *Spodoptera exiguum* (Hübner, [1808])

175. *Tyta luctuosa* ([Denis & Schiffermüller], 1775)

176. *Ulochlaena hirta* (Hübner, [1813])

Locally common. Autumnal.

177. *Xylena exsoleta* (Linnaeus, 1758)

Vernal, Autumnal.

Pyralidae

Although this family is well represented in Turkey, the current short list is the preliminary one.

178. *Chrysocrambus linetellus* (Fabricius, 1781)

179. *Eurrhypis cacuminalis* (Eversmann, 1843)

Diurnal, rare in spring.

180. *Glaucocharis euchromiella* (Ragonot, 1895)

181. *Loxostege peltaloides* (Rebel, 1932) (Fig. 26)

Autumnal.

182. *Metacrambus caretellus* (Zeller, 1847)

183. *Psammotis pulveralis* (Hübner, 1796)

184. *Pyrausta purpuralis* (Linnaeus, 1758)

185. *Synaphe consecratalis* (Lederer, 1855)

186. *Synaphe moldavica* (Esper, [1789])

187. *Udea ferrugalis* (Hübner, 1796)

188. *Udea praepetalis* (Lederer, 1869)

Scythrididae

189. *Scythris (scopolella-gr.) flabella* (Mann, 1861)

Sphingidae

190. *Hemaris (Cochrania) croatica* (Esper, [1779])

Diurnal.

191. *Hyles hippophaes* (Esper, [1789])

Tortricidae

192. *Cnephasia (s.str.) grandis* (Ostheder, 1938)

193. *Eugnosta magnificana* (Rebel, 1914)

194. *Fulvoclysia subdolana* (Kennel, 1901)

Zygaenidae

195. *Zygaena (Agrumenia) carniolica* (Scopoli, 1763) Locally common.

196. *Zygaena (Mesembrynus) tamara* Christoph, 1889 Locally common.

Selected images on the earlier observations in nature

Figs. 1, 2 – Caterpillars of *Polygonia egea* on *Parietaria judaica*. From Erek Mt 2200m on 2-4 6 2008 M. Kemal (Cesa)



Figs. 3, 4 – Prepupal stage on 10 6 2008 (left); pupal skin and emerged butterfly on 22 6 2008, M. Kemal (Cesa)



Figs. 5,6 - *Carcharodus alceae*, from Sarmaç (Erek Mt.) 2100m, on 5 6 2007 (left); *Pyrgus sidae* from Sarmaç (Erek Mt.) on 2 6 2011 (right), M. Kemal (Cesa)



Figs. 7, 8 - *Lycaena asabinus*, a male feeding on the flower of *Senecio* (Asteraceae). Erek Mt., Sarmaç 2100m on 26 2011 (left); *Lycaena kefersteini* (male), feeding on the same flower, at the same place and date (right), M. Kemal (Cesa)



Figs. 9, 10 - *Lycaena kurdistanica*, a male feeding on the flower of *Ornithogalum* (Liliaceae). Erek Mt., Kevenli 2000m, on 8 6 2011 (left); a female of the same species, at the same place and date (right), M. Kemal (Cesa)



Figs. 11, 12 - *Lycaena phlaeas*, a feeding butterfly on the flower of *Senecio* (Asteraceae). Erek Mt., Sarmaç 2100m on 26 2011 (left); a male of *Rubrapterus bavius* ssp. *vanicola*, feeding on the flowers of *Aethionema* (Brassicaceae), Erek Mt., Hoşpurak 2000m, on 13 6 2009 (right), M. Kemal (Cesa)



Figs. 13, 14 - *Anthocharis damone* (male), from Sarmaç (Erek Mt.) 2100m, on 16 6 2007 (left); *Anthocharis grueneri* (female) from Sarmaç (Erek Mt.) on 2 6 2011 (right), M. Kemal (Cesa)



Figs. 15, 16 - *Lasiommata maera*, at rest. Erek Mt., Hoşpurak 2000m, on 13 6 2009 (left); *Aplocera annexata*, at rest. From same place and date (right), M. Kemal (Cesa)

Images of recent observations in nature



Fig. 17 – Habitat of the autumnal moths. Erek Mt. 2200m, on 19 October 2017 evening hours, M. Kemal (Cesa)



Figs. 18, 19 - *Larentia clavaria*, nocturnal. Erek Mt., Sarmaç 2150m (65Kl3), on 20 10 2017 (left); *Scotopteryx kuznetzovi*, nocturnal. Erek Mt., Sarmaç 2150m (65Kl3), on 20 10 2017 (right), M. Kemal (Cesa)



Figs. 20, 21 - *Malacosoma castrensis* overwintering eggs cluster. Erek Mt. Sarmaç 2150m (65Kl3), on 17 10 2017 (left); *Agrochola egorovi*, nocturnal. Erek Mt., Sarmaç 2150m (65Kl3), on 21 10 2017 (right), M. Kemal (Cesa)




Figs. 22, 23 – *Dichagyris amoena*, nocturnal. Erek Mt., Sarmaç 2150m (65Kl3), on 21 10 2017 (left); *Heptapotamia eustrati*, nocturnal. From same place and date (right), M. Kemal (Cesa)




Figs. 24, 25 – *Margelana flavidior*, nocturnal. Erek Mt., Sarmaç 2150m (65Kl3), on 21 10 2017 (left); *Polymixis perchrysa*, nocturnal. From same place and date (right), M. Kemal (Cesa)




Fig. 26 – *Loxostege peltaloides*, nocturnal. Erek Mt., Sarmaç 2150m (65Kl3), on 21 10 2017 , M. Kemal (Cesa)

Images of the specimens with their genitalia and tympanal organs

Figs. 27, 28 – *Scotopteryx kuznetzovi*. Upperside of male (left), and its genitalia. GP2750♂, Erek Mt., Sarmaç 2150m (65K13), on 20 10 2017 (right), M. Kemal (Cesa)



Figs. 29, 30 - *Scotopteryx kuznetzovi*. Tympanal organ of the male (before and after preparation. GP2750, M. Kemal (Cesa)



Figs. 31, 32 – *Agrochola egorovi*. Male genitalia, with removed aedeagus with enlarged parts and abdominal skin. GP2771. Erek Mt. Sarmaç 2150m on 20 10 2017, M. Kemal (Cesa)



Fig. 33 - *Polymixis csorbagabori*, Upperside of male. GP2753. Erek Mt., Sarmaç 2150m, 21 10 2017, M. Kemal (Cesa)




Fig. 34 – Male genitalia of *Polymixis csorbagabori*, GP2753. Aedeagus removed, two parts of the genitalia enlarged. Erek Mt. Sarmaç 2150m, M. Kemal (Cesa)




Fig. 35 – *Polymixis perchrysa*. Male genitalia. GP2756. Erek Mt. Sarmaç 2150m (65Kl3), on 21 10 2017, M. Kemal (Cesa)



Figs. 36, 37 – *Polymixis rosinae*. Upperside of male (left), and male genitalia. GP2755. Erek Mt., Sarmaç 2150m, 21 10 2017, M. Kemal (Cesa)

References

Boursin,C., 1944, Contribution a l'étude de la faune du Caucase et de l'Arménie. *Revue fr. Ent.* 10: 75-84, Pls.

Hacker, H. & L.Ronkay, 1992, Das Genus Polymixis Hübner,[1820] mit Beschreibung neuer Taxa und Festlegung neuer Stati (Lepidoptera, Noctuidae). *Esperiana* 3: 473-496, figs.

Kayci, L., 2007, Studies on the fauna and ecology of Papilionoidea and Hesperioidea in Erek Mountain (Van). *Priamus Suppl.* 6: 1-47.

Kemal, M., 2009, Biodiversity in Erek Mountain and its rarest butterfly *Callophrys mystaphia* (Lepidoptera, Lycaenidae). MPEG2 file. Duration 8 min. 04 sec.
<http://www.archive.org/details/BiodiversityInErekMountainAndItsRarestButterflyCallophrysMystaphiatr> [subtitle in Turkish]
<http://www.archive.org/details/BiodiversityInErekMountainAndItsRarestButterflyCallophrysMystaphiaeg> [subtitle in English]

Kemal, M. & A.Ö.Koçak, 2015, Pictorial notes on the early stages of *Heptapotamia eustratii* Alph. in East Turkey (Lepidoptera, Noctuidae). *Cesa News* 112: 1-6, 14 figs.

Kemal,M. & A.Ö.Koçak, 2017, On the Lepidoptera fauna of a single locality, Saklivadi (Van Province, East Turkey). *Priamus* 15 (1): 1-80, 101 figs., 45 maps, 1 Tab.

Koçak,A.Ö. & M. Kemal, 2012, List of two thousand species of pterygot insects in Van Province (East Turkey) (Results of the entomofauna project of Turkey – 7). *Cesa News* 81: 2-86, 37 figs.

Rajaei,H.S. & D.Stünning, 2012. *Scotopteryx kuznetzovi* (Wardikian,1957) (Lepidoptera, Geometridae, Larentiinae), a new species for the fauna of Iran and Turkey. *Bonn. Zool. Bull.* 61 (1): 135-139, figs.

Ronkay,L., 1984, Notes on the genus *Agrochola* Hübner,1821 (Lepidoptera, Noctuidae). Part II. *Acta zool. hung.* 30 (1-2): 179-187, figs.

Cesa News

Cesa News is a peer reviewed online serial of the Centre for Entomological Studies Ankara (Cesa), established in 2008. It appears at irregular intervals as PDF format and includes original articles of the research workers of the Centre, regarding on various subjects on Entomology. The publication language is English or Turkish. Cesa News is archived online at "Internet Archive" since 2011, in accordance with the publication rules of the ICZN.

Cesa News is an open-access serial, distributed under the terms of the "*Creative Commons Attribution License*", which permits free use, and distribution in any medium, provided the original author(s) and source are credited.

Centre for Entomological Studies Ankara



(A scientific Consortium)
(co-operation of research workers for pure-scientific, not commercial purpose)

Web Page of the Cesa: <http://www.cesa-tr.org/> - Digital Library of the Cesa: <https://archive.org/details/@cesa>

Scientific Serials: Priamus & Priamus Supplement (print and online versions) (ISSN 1015-8243)², Miscellaneous Papers (print and online versions) (ISSN 1015-8235)³, Memoirs (print and online versions) (ISSN-8227)⁴ DVD Films⁵, Iconographia Insectorum⁶ (online), Cesa Publications on African Lepidoptera (online)⁷, Cesa News (online)⁸, Cesa Books (online)⁹

Owners / Sahipleri - Editors / Yayıncılar: Prof. em. Dr. Ahmet Ömer Koçak (c/o Yüzüncü Yıl University, Van, Turkey), Asst. Prof. Dr. Muhabbet Kemal Koçak (c/o Yüzüncü Yıl University, Van, Turkey).

Editorial Board of all Scientific Serials of the CESA / Bütün Bilimsel Yayınların Yayın Kurulu: Insecta, taxonomy, nomenclature, ecology, faunistics: Prof. Dr. Ahmet Ömer Koçak (Yüzüncü Yıl Üniversitesi, Turkey), Asst. Prof. Dr. Muhabbet Kemal Koçak (Yüzüncü Yıl University, Turkey).

Chief referees of all Scientific Serials of the CESA: Prof. em. Dr. Ahmet Ömer Koçak & Asst. Prof. Dr. Muhabbet Kemal Koçak: Insecta, taxonomy, nomenclature, fauna, ecology, catalogues, checklists of the Old World.

Expert referees according to the subject areas: Dr. Peter Huemer (Austria): Gelechiidae, and some Microlepidoptera groups in Palaearctic (Lepidoptera). Dr. J. B. Heppner (U.S.A.): Microlepidoptera of Nearctic and Neotropical. Dr. G. Baldizzone (Italy): Coleophoridae (Lepidoptera). Dr. V. Korneyev (Ukraine): Tephritidae, Pyrgotidae, Ulidiidae (Diptera). Prof. Dr. Y.G. Verves (Ukraine): Sarcophagidae (Diptera). Dr. Daniel Burckhardt (Switzerland): Psyllidae (Homoptera). Prof. Dr. E. Heiss (Austria): Hemiptera. Dr. R. Ehrmann (Germany): Mantodea. Prof. Dr. Mustafa Ünal (Bolu, Turkey): Orthoptera. Prof. Dr. Hüseyin Özdişmen (Turkey): Coleoptera. Prof. Dr. Suat Kiyak (Turkey): Hemiptera, Hymenoptera (Cynipidae).

Plant taxonomy, flora and vegetation: Assoc. Prof. Dr. Murat Ünal, Asst. Prof. Dr. Mesut Pınar, (Yüzüncü Yıl University, Van, Turkey).

Editorial policy: The submitted manuscript is evaluated by the Chief Editor and Referee. In case of need, the manuscript is sent to expert referees according to the subject areas.

ALL RIGHTS RESERVED

Correspondences should be addressed to: Prof. em. Dr. Ahmet Ömer Koçak, c/o Yüzüncü Yıl University, Fen Fakültesi, Biyoloji Bölümü, Kampus, Van / Turkey. - e-mail: cesa_tr@yahoo.com.tr

All the serials of the Cesa are archived regularly by the Internet Archive

(300 Funston Ave., San Francisco, CA 94118, U.S.A.),

in accordance with the rules of the International Codes of Zoological Nomenclature (ICZN)

<https://archive.org/>

also recorded regularly by the Zoological Record,

Thomson Reuters, Enterprise House, Innovation Way, Heslington, York, YO10 5NQ, United Kingdom

<https://clarivate.com/>

² <http://www.cesa-tr.org/Priamus.htm> <http://www.cesa-tr.org/Pri.htm>

³ <http://www.cesa-tr.org/Miscell.htm>

⁴ <http://www.cesa-tr.org/Memoirs.htm>

⁵ <http://www.cesa-tr.org/CDF.htm>

⁶ <http://www.cesa-tr.org/Icon.htm>

⁷ <http://www.cesa-tr.org/CPAL.htm>

⁸ <http://www.cesa-tr.org/Cesanews.htm>

⁹ <http://www.cesa-tr.org/Cesabooks.htm>